

NEW FLAVOR CHANGING INTERACTIONS IN EXTENDED GAUGE MODELS [†]

ENRICO NARDI

*Randall Laboratory of Physics, University of Michigan
Ann Arbor, MI 48109-1120, U.S.A.*

ABSTRACT

A new class of flavor changing (FC) neutral interactions can arise in models based on extended gauge groups (rank >4) when new charged fermions are present together with new neutral gauge bosons. In some cases the FC couplings to a new Z_1 are expected to be sizeable, implying that the Z_1 mass must be large enough as to explain the observed suppression of FC transitions. Concentrating on E_6 models and assuming for the FC parameters a theoretically natural range of values, I show that in most cases the presence of a Z_1 much lighter than 1 TeV is unlikely.

PACS number(s): 13.10.+q, 12.10.Dm, 12.15.Cc, 14.60.Jj

[†] Talk given at the 7th Meeting of the American Physical Society, DPF92, 10-14 November 1992, Fermi National Accelerator Laboratory, Batavia, Illinois.

E-mail: nardi@umiphys.bitnet

1. Introduction

The Standard Model (SM) of the electroweak interactions has achieved a tremendous success in describing the experimental data within the range of energies available today. The present theory also accommodates in a satisfactory way the whole spectrum of known particles, and only two states, the top-quark and the Higgs boson, necessary for the consistency of the model, have not been discovered yet. In addition a large set of experimental limits on rare processes are explained in a satisfactory way, since the model includes the Glashow-Iliopoulos-Maiani mechanism for suppressing flavor changing neutral currents (FCNC) in the quark sector, and strictly forbids lepton flavor violating (LFV) currents. All these features are quite peculiar to the SM, and in general most of its possible extensions predict a larger spectrum of states as well as larger rates for FCNC processes.

Here I will analyze a new class of FCNC interactions that are generally present in most of the extensions of the SM which predict one (or more) additional neutral gauge boson Z_1 together with new charged fermions, and that are induced by Z_1 interactions. In some cases the FC couplings to the Z_1 are not suppressed, and when one considers the current limits on the corresponding FC transitions this points toward a rather heavy Z_1 ¹. In general the standard Z_0 is expected to be mixed with the Z_1 . The resulting mass eigenstate will then acquire new FC couplings proportional to the Z_0 – Z_1 mixing angle¹. However in this short presentation I will neglect these additional FC effects, since due to the tight limits implied for the Z_0 – Z_1 mixing by low energy NC and LEP data ($\phi_{Z_0-Z_1} \lesssim 0.02$ ^{2,3}) they turn out to be less important than the effects due to Z_1 exchange¹. In order to illustrate the power of the constraints which can be derived from the limits on FCNC, I will apply them to a class of E_6 grand unified theories (GUTs), and I will show that once a natural range of values for the new FC parameters is assumed, the non-observation of the decay $\mu \rightarrow eee$ tightly constraints the mass of the Z_1 .

2. Formalism

Once a low energy gauge group of the form $[SU(2)_L \times U(1)_Y \times SU(3)_C] \times U_1(1)$ is assumed, the neutral current Lagrangian in the gauge basis reads

$$-\mathcal{L}_{\text{NC}} = eJ_{\text{em}}^\mu A_\mu + g_0 J_0^\mu Z_{0\mu} + g_1 J_1^\mu Z_{1\mu}. \quad (1)$$

The SM neutral gauge boson Z_0 couples with strength $g_0 = (4\sqrt{2}G_F M_{Z_0}^2)^{1/2}$ to the usual combination of the neutral isospin and electromagnetic currents $J_0^\mu = J_3^\mu - \sin^2 \theta_W J_{\text{em}}^\mu$. Assuming that the new $U_1(1)$ originates from a GUT based on a *simple* group, and normalizing the new generator Q_1 to the hypercharge axis, then the Z_1

couples to the new J_1 current with strength $g_1 \simeq g_0 \sin \theta_W$. To ensure the absence of anomalies for the new gauge current J_1 , *new* fermions must be present in addition to the standard 15 *known* fermions per generation. Here I will assume that some of the additional new fermions are electrically charged, and that they are mixed with the known states. Each of the conventional *light* fermion mass eigenstate then corresponds to a superposition of the known states and the new states. Conservation of the electric and color charges forbids a mixing between gauge eigenstates with different $U(1)_{\text{em}}$ and $SU(3)_c$ quantum numbers, implying in turn that the corresponding currents are not modified by the presence of the new states. In contrast the neutral isospin generator T_3 and the new generator Q_1 are spontaneously broken, and a mixing between states with different t_3 and q_1 eigenvalues is allowed. This will affect the J_3 and J_1 currents² and in turn the couplings of the light mass eigenstates to the Z_0 and Z_1 . In the gauge currents chirality is conserved too, and it is then convenient to group the fermions with the same electric charge and chirality $\alpha = L, R$ in a vector of the known ($\mathcal{K}$) and new ($\mathcal{N}$) gauge eigenstates² $\Psi_\alpha^o = (\Psi_\mathcal{K}^o, \Psi_\mathcal{N}^o)^T_\alpha$. This vector is related to the corresponding vector of the light (l) and heavy (h) mass eigenstates $\Psi_\alpha = (\Psi_l, \Psi_h)^T_\alpha$ through a unitary transformation

$$\begin{pmatrix} \Psi_\mathcal{K}^o \\ \Psi_\mathcal{N}^o \end{pmatrix}_\alpha = U_\alpha \begin{pmatrix} \Psi_l \\ \Psi_h \end{pmatrix}_\alpha \quad \text{where} \quad U_\alpha = \begin{pmatrix} A & G \\ F & H \end{pmatrix}_\alpha, \quad \alpha = L, R. \quad (2)$$

In terms of the fermion mass eigenstates, the neutral current corresponding to a (broken) generator $\mathcal{Q} = T_3, Q_1$ now reads

$$J_\mathcal{Q}^\mu = \sum_{\alpha=L,R} \bar{\Psi}_\alpha \gamma^\mu U_\alpha^\dagger \mathcal{Q}_\alpha U_\alpha \Psi_\alpha, \quad (3)$$

where $\mathcal{Q}_\alpha$ represents a generic diagonal matrix of the charges $q_\alpha = t_3(f_\alpha), q_1(f_\alpha)$ for the chiral fermion f_α . Since we are interested in the indirect effects of fermion mixings in the couplings of the light mass eigenstates, we have to project $J_\mathcal{Q}^\mu$ onto the light components Ψ_l . In the particularly simple case when the mixing is with only one type of new fermions with the same $q_\alpha^\mathcal{N}$ charges, by means of the unitarity of U_α we easily obtain²

$$J_{l\mathcal{Q}}^\mu = \sum_{\alpha=L,R} \bar{\Psi}_{l\alpha} \gamma^\mu [q_\alpha^\mathcal{K} I + (q_\alpha^\mathcal{N} - q_\alpha^\mathcal{K}) F_\alpha^\dagger F_\alpha] \Psi_{l\alpha}. \quad (4)$$

In Eq. (4) $q_\alpha^\mathcal{K} I$ represents the coupling of a particular light fermion in the absence of mixing effects, while the second term accounts for the modifications due to fermion mixings. The matrix $F_\alpha^\dagger F_\alpha$ is in general not diagonal, and clearly whenever the coefficient $(q_\alpha^\mathcal{N} - q_\alpha^\mathcal{K})$ is nonvanishing, the off diagonal terms will induce FCNC. We can distinguish two cases. When the mixing violates weak-isospin ($t_3(f_\alpha^\mathcal{N}) \neq t_3(f_\alpha^\mathcal{K})$) the J_0 current is affected, and the Z_0 interactions will be flavor changing. However an analysis of the fermion mass matrix¹ shows that the isospin-violating fermion

mixings, which are generated via $\Delta t_3 = \frac{1}{2}$ off-diagonal mass terms, are suppressed as the ratio of the heavy to light masses. Hence the corresponding FC terms $(F_\alpha^\dagger F_\alpha)_{i \neq j}$ are predicted to be quite small, and hence there is no theoretical conflict with the experimental limits⁴. In contrast $\Delta t_3 = 0$ mass terms induce mixings that do not violate weak isospin, and it turns out that in this case no suppression factors are present¹. Since in this case $t_3(f_\alpha^\mathcal{N}) = t_3(f_\alpha^\mathcal{K})$, clearly the J_0 current is not affected and remains flavor-diagonal. However, in general we still have $q_1(f_\alpha^\mathcal{N}) \neq q_1(f_\alpha^\mathcal{K})$, and then the isospin-conserving mixings can affect the J_1 current, inducing sizeable FC couplings to the Z_1 . This constrains the Z_1 mass to be sufficiently heavy for suppressing at low energy the FC transitions via propagator effects. Similarly a possible Z_0 – Z_1 mixing would induce additional FC contributions to the vertices of the Z mass eigenstate, therefore the mixing cannot be too large so as to conflict with the experimental bounds¹.

3. Constraints on E_6 models from $\mu \rightarrow eee$.

E_6 GUTs are well known examples of theories where additional fermions and new neutral gauge bosons are simultaneously present, giving rise to the kind of effects which I have discussed. For a general breaking of E_6 (rank 6) to the SM (rank 4) it is possible to define a whole class of Z_1 bosons corresponding to a linear combination of the two additional Cartan generators⁵. I will parametrize this combination in terms of an angle β . Fermions are assigned to the fundamental **27** representation of the group which contains 12 additional states for each generation, among which we have a vector doublet of new leptons $(N E^-)_L^T, (E^+ N^c)_L^T$. Non-diagonal mass terms with the standard $(\nu e^-)_L^T$ and e_L^c leptons will give rise respectively to $\Delta t_3 = 0$ and $\Delta t_3 = \frac{1}{2}$ mixings, and in particular will induce LFV left (L) and right (R) chiral couplings between the first and second generation, allowing for the decay $\mu \rightarrow eee$ ¹. A very stringent experimental limit exists for this decay mode⁶: $Br(\mu^+ \rightarrow e^+ e^+ e^-) < 1.0 \cdot 10^{-12}$. In order to derive bounds for the Z_1 mass from this result I will conservatively neglect the LFV couplings in the R sector, and assume that the only source of the LFV interaction comes from the $\Delta t_3 = 0$ mixing in the L sector. I will also assume that the LFV term $\mathcal{F}_{e\mu} \equiv (F_L^\dagger F_L)_{e\mu}$ lies in the ‘natural’ range 10^{-2} – 10^{-3} . This assumption relies on the observation that the CKM mixings, which are also isospin-conserving, are numerically $> 10^{-3}$ and that the mixing between the first and second generation is particularly large. The limits on the Z_1 mass obtained in this way¹ are depicted in Fig.1 as a function of the parameter β that defines the particular E_6 boson. These limits are indeed very strong, however, since they depend on a specific assumption for the numerical value of the LFV coupling,

Figure 1: Limits on M_{Z_1} from $\mu \rightarrow eee$ for a general neutral gauge boson from E_6 , as a function of $\sin\beta$. The values of $\sin\beta$ corresponding to the particularly interesting Z_η , Z_χ and Z_ψ bosons, are also shown at the top of the figure. The limits are given for the two different values of the LFV term $\mathcal{F}_{e\mu} = 10^{-2}$ and 10^{-3} .

they cannot replace the direct⁷ bounds or other more model independent indirect limits^{2,3}.

4. References

1. E. Nardi, Preprint UM-TH-92-19.
2. E. Nardi, E. Roulet and D. Tommasini, *Phys. Rev.* **D46**, (1992) 3040.
3. G. Altarelli *et.al.*, *Phys. Lett.* **B263**, (1991) 459;
M.C. Gonzalez García and J.W.F. Valle; *Phys. Lett.* **B259**, (1991) 365;
F.del Aguila, J.M. Moreno and M. Quirós, *Nucl. Phys.* **B361**, (1991) 45;
F. del Aguila, W. Hollik, J.M. Moreno, M. Quirós, *Nucl. Phys.* **B372**, (1992) 3; *ibid.* **B372** (1992) 3;
P. Langacker and M. Luo, *Phys. Rev.* **D45**, (1992) 278;
J. Layssac, F.M. Renard and C. Verzegnassi, *Z. Phys.* **C53**, (1992) 97.
4. P. Langacker and D. London, *Phys. Rev.* **D38**, (1988) 886.
5. J.L. Hewett and T.G. Rizzo, *Phys. Rep.* **183**, (1989) 195.
6. SINDRUM collaboration, U. Bellgardt *et.al.*, *Nucl. Phys.* **B299**, (1988) 1.
7. CDF Collaboration, F. Abe *et.al.*, *Phys. Rev. Lett.* **68**, (1992) 1463.